arXiv:1611.05610v1 [math.AG] 17 Nov 2016

ON REAL ANTI-BICANONICAL CURVES WITH ONE DOUBLE POINT ON THE 4-TH REAL HIRZEBRUCH SURFACE. II

SACHIKO SAITO

Abstract. A real 2-elementary K3 surfaces of type $((3,1,1),-\mathrm{id})$ yields a real anti-bicanonical curve $s \cup A_1'$ (disjoint union) on the 4-th real Hirzebruch surface $\mathbb{F}_4$ where s is the exceptional section of $\mathbb{F}_4$ and the real curve A_1' has one real double point. We give a criterion (see Lemma 2.3) which determines whether the real double point is degenerate or not. One direction of the assertion of Lemma 2.3 has already been proved in Lemma 4.6 in the preceding paper [8]. In this paper we prove the inverse direction.

Contents

1. Introduction: Review of real 2-elementary K3 surfaces 1
1.1. Real 2-elementary K3 surfaces 1
1.2. Integral involutions of $\mathbb{L}_{K3}$ of type (S,θ) 3
1.3. Period domains 4
1.4. $(\mathcal{D}\mathbb{R})$-nondegenerate marked real 2-elementary K3 surfaces 5
2. Real 2-elementary K3 surfaces of type $((3,1,1),-\mathrm{id})$ 5
References 9

1. Introduction: Review of real 2-elementary K3 surfaces

1.1. **Real 2-elementary K3 surfaces.** A K3 surface X with a non-symplectic holomorphic involution τ is often called a 2-*elementary K3 surface* (X,τ). Note that every K3 surface with a non-symplectic holomorphic involution is algebraic.

Definition 1.1 (Real 2-elementary K3 surface)**.** A triple (X,τ,φ) is called a *real K3 surface with non-symplectic holomorphic involution* (or *real* 2*-elementary K3 surface*) if
(1) (X,τ) is a K3 surface X with a non-symplectic holomorphic involution τ.
(2) φ is an anti-holomorphic involution on X.
(3) $\varphi\circ\tau=\tau\circ\varphi$.

For a 2-elementary K3 surface (X,τ), generally, we denote by

$$H_{2+}(X,\mathbb{Z})$$

the fixed part of $\tau_*: H_2(X,\mathbb{Z})\to H_2(X,\mathbb{Z})$. It is well-known that $H_2(X,\mathbb{Z})$ is an even unimodular lattice of signature $(3,19)$. Such lattices are all isometric.

$H_{2+}(X,\mathbb{Z})$ is a primitive hyperbolic 2-elementary sublattice of $H_2(X,\mathbb{Z})$. Note that $H_{2+}(X,\mathbb{Z})$ is contained in $\mathrm{Pic}(X)$, where $\mathrm{Pic}(X)$ is the Picard lattice of X.

---

partially supported by JSPS Grant-in-Aid for Challenging Exploratory Research 25610001 (2013/4 — 2016/3).
2010 AMS Mathematics Subject Classification: 14J28, 14P25, 14J10.

We fix an even unimodular lattice $\mathbb{L}_{K3}$ of signature $(3,19)$. Let S $(\subset \mathbb{L}_{K3})$ be a primitive hyperbolic 2-elementary sublattice of $\mathbb{L}_{K3}$.

Definition 1.2. We set $r(S) := \operatorname{rank} S$. The non-negative integer $a(S)$ is defined by $S^*/S \cong (\mathbb{Z}/2\mathbb{Z})^{a(S)}$. We define the "parity"

$$\delta(S) := \begin{cases} 0 & \text{if } z \cdot \sigma(z) \equiv 0 \bmod 2 \ \ (\forall z \in \mathbb{L}_{K3}) \\ 1 & \text{otherwise,} \end{cases}$$

where $\sigma : \mathbb{L}_{K3} \to \mathbb{L}_{K3}$ is the unique integral involution whose fixed part is S.

It is known that the triplet $(r(S), a(S), \delta(S))$ determines the isometry class of the lattice S ([5]).

If S and S' are isometric primitive hyperbolic 2-elementary sublattices of the K3 lattice $\mathbb{L}_{K3}$, then there exists an ambient automorphism f of $\mathbb{L}_{K3}$ such that $f(S') = S$ ([1], [4]). Hence, if (X, τ) and (X', τ') are two 2-elementary K3 surfaces, $\alpha : H_2(X, \mathbb{Z}) \to \mathbb{L}_{K3}$ is an isometry (so-called "marking" as below), and $H_{2+}(X, \mathbb{Z})$ is isometric to $H_{2+}(X', \mathbb{Z})$; then there exists an isometry $\alpha' : H_2(X', \mathbb{Z}) \to \mathbb{L}_{K3}$ such that $\alpha'(H_{2+}(X', \mathbb{Z})) = \alpha(H_{2+}(X, \mathbb{Z}))$. ([8])

We additionally fix a half cone (so-called "positive cone", see [6])

$$V^+(S)$$

of the cone

$$V(S) := \{x \in S \otimes \mathbb{R} \mid x^2 > 0\}.$$

We moreover fix a fundamental subdivision

$$\Delta(S) = \Delta(S)_+ \cup -\Delta(S)_+$$

of all elements with square -2 in S. This is equivalent to fixing a fundamental chamber

$$\mathcal{M} \ \ (\subset V^+(S))$$

for the group $W^{(-2)}(S)$ generated by reflections in all elements with square (-2) in S.

Note that $\mathcal{M}$ and $\Delta(S)_+$ define each other by the condition $\mathcal{M} \cdot \Delta(S)_+ \geq 0$.

Let θ be an integral involution of S.

Definition 1.3. We say (X, τ, φ) is a *real 2-elementary K3 surface of type* (S, θ) if there exists an isometry $\alpha : H_2(X, \mathbb{Z}) \cong \mathbb{L}_{K3}$ such that $\alpha(H_{2+}(X, \mathbb{Z})) = S$ and the following diagram commutes:

$$\begin{array}{ccc} H_{2+}(X, \mathbb{Z}) & \xrightarrow{\alpha} & S \\ {\scriptstyle \varphi_*}\downarrow & & \downarrow{\scriptstyle \theta} \\ H_{2+}(X, \mathbb{Z}) & \xrightarrow{\alpha} & S. \end{array}$$

Definition 1.4 (marked real 2-elementary K3 surfaces)**.** A *marked real 2-elementary K3 surface of type* (S, θ) is defined to be a pair $((X, \tau, \varphi),\ \alpha)$ of a real 2-elementary K3 surface (X, τ, φ) of type (S, θ) (Definition 1.3) and an isometry, which is called *marking*,

$$\alpha : H_2(X, \mathbb{Z}) \cong \mathbb{L}_{K3}$$

such that

- $\alpha(H_{2+}(X, \mathbb{Z})) = S$,
- $\alpha \circ \varphi_* = \theta \circ \alpha$ on $H_{2+}(X, \mathbb{Z})$,
- $\alpha_{\mathbb{R}}^{-1}(V^+(S))$ contains a hyperplane section of X, and

- the set $\alpha^{-1}(\Delta(S)_+)$ contains only effective classes of X.

Here we say a class in $H_2(X,\mathbb{Z})$ is *effective* if it is represented by an effective divisor of X. (See, for example, [2].)

Remark 1.5. Note that ([6]) for any (X,τ), we can take α such that $\alpha_{\mathbb{R}}^{-1}(V^+(S))$ contains a hyperplane section of X.

1.2. Integral involutions of $\mathbb{L}_{K3}$ of type (S,θ).

Definition 1.6 (Integral involution ψ of $\mathbb{L}_{K3}$ of type (S,θ))**.** Let S be a hyperbolic 2-elementary sublattice of $\mathbb{L}_{K3}$ (as above), $\theta : S \to S$ be an integral involution. Let $\psi : \mathbb{L}_{K3} \to \mathbb{L}_{K3}$ be an integral involution of the lattice $\mathbb{L}_{K3}$ such that the following diagram commutes:

$$\begin{array}{ccc} S & \subset & \mathbb{L}_{K3} \\ \theta \downarrow & & \downarrow \psi \\ S & \subset & \mathbb{L}_{K3}. \end{array}$$

We say such a pair $(\mathbb{L}_{K3},\psi)$ (or ψ) an *integral involution of* $\mathbb{L}_{K3}$ *of type* (S,θ).

Definition 1.7 (the associated integral involution)**.** Let $((X,\tau,\varphi),\ \alpha)$ be a marked real 2-elementary K3 surface of type (S,θ). If we set

$$\psi := \alpha \circ \varphi_* \circ \alpha^{-1} : \mathbb{L}_{K3} \to \mathbb{L}_{K3},$$

then we have $\psi(S) = S$, and $\psi(x) = \theta(x)$ for every $x \in S$ because $\alpha^{-1}(x) \in H_{2+}(X,\mathbb{Z})$. Hence, $(\mathbb{L}_{K3},\psi)$ is an integral involution of $\mathbb{L}_{K3}$ of type (S,θ).

We call the integral involution $(\mathbb{L}_{K3},\psi)$ of type (S,θ) *the associated integral involution* with a marked real 2-elementary K3 surface $((X,\tau,\varphi),\ \alpha)$ of type (S,θ) if the following diagram commutes:

$$\begin{array}{ccc} H_2(X,\mathbb{Z}) & \xrightarrow{\alpha} & \mathbb{L}_{K3} \\ {\scriptstyle\varphi_*}\big\downarrow & & \big\downarrow{\scriptstyle\psi} \\ H_2(X,\mathbb{Z}) & \xrightarrow{\alpha} & \mathbb{L}_{K3}. \end{array}$$

Definition 1.8. Let $\Delta(S,L)^{(-4)}$ be the set of all elements δ_1 in S such that $\delta_1^2 = -4$ and there exists $\delta_2 \in S_L^\perp$ such that $(\delta_2)^2 = -4$ and $\delta = (\delta_1+\delta_2)/2 \in L$. Let $W^{(-4)}(S,L)$ $(\subset O(S))$ be the group generated by reflections in all elements in $\Delta(S,L)^{(-4)}$, and $W^{(-4)}(S,L)_{\mathcal{M}}$ be the stabilizer subgroup of $\mathcal{M}$ in $W^{(-4)}(S,L)$. We define the subgroup G to be generated by reflections s_{δ_1} in all elements $\delta_1 \in \Delta(S,L)^{(-4)}$ which are contained either in S_+ or in S_- and satisfy $(s_{\delta_1})_{\mathbb{R}}(\mathcal{M}) = \mathcal{M}$. G is a subgroup of $W^{(-4)}(S,L)_{\mathcal{M}}$.

Definition 1.9 (Isometries with respect to the group G)**.** Let $(\mathbb{L}_{K3},\psi_1)$ and $(\mathbb{L}_{K3},\psi_2)$ be two integral involutions of $\mathbb{L}_{K3}$ of type (S,θ). An *isometry with respect to the group* G from $(\mathbb{L}_{K3},\psi_1)$ to $(\mathbb{L}_{K3},\psi_2)$ means an isometry $f : \mathbb{L}_{K3} \to \mathbb{L}_{K3}$ such that $f(S) = S$, $f|_S \in G$, and the following diagram commutes:

$$\begin{array}{ccc} \mathbb{L}_{K3} & \xrightarrow{f} & \mathbb{L}_{K3} \\ \big\downarrow{\scriptstyle\psi_1} & & {\scriptstyle\psi_2}\big\downarrow \\ \mathbb{L}_{K3} & \xrightarrow{f} & \mathbb{L}_{K3}. \end{array}$$

Definition 1.10. We say two integral involutions $(\mathbb{L}_{K3}, \psi_1)$ and $(\mathbb{L}_{K3}, \psi_2)$ of type (S, θ) are *isometric with respect to the group G* if there exists an isometry with respect to the group G from $(\mathbb{L}_{K3}, \psi_1)$ to $(\mathbb{L}_{K3}, \psi_2)$.

By an *automorphism* of an integral involution $(\mathbb{L}_{K3}, \psi)$ of type (S, θ) *with respect to the group G* we mean an isometry with respect to the group G from $(\mathbb{L}_{K3}, \psi)$ to itself. Namely, an isometry $f : \mathbb{L}_{K3} \to \mathbb{L}_{K3}$ which satisfies that $\psi \circ f = f \circ \psi,\ f(S) = S$ and $f|_S \in G$.

Definition 1.11 (analytic isomorphisms with respect to G)**.** Two marked real 2-elementary K3 surfaces $((X, \tau, \varphi),\ \alpha)$ and $((X', \tau', \varphi'), \alpha')$ of type (S, θ) are *analytically isomorphic with respect to the group G* if there exists an analytic isomorphism $f : X \to X'$ such that $f \circ \tau = \tau' \circ f$, $f \circ \varphi = \varphi' \circ f$ and $\alpha' \circ f_* \circ \alpha^{-1} | S \in G$.

1.3. **Period domains.** Let us **fix** an integral involution

$$(\mathbb{L}_{K3}, \psi)$$

of type (S, θ) through this subsection.

We follow the formulations of period domains of marked real 2-elementary K3 surfaces in Itenberg [3] and Nikulin-Saito [6].

We set

$$\Omega_\psi := \{\omega\ (\neq 0) \in \mathbb{L}_{K3} \otimes \mathbb{C} \mid \omega \cdot \omega = 0,\ \omega \cdot \overline{\omega} > 0,\ \omega \cdot S = 0,\ \psi_{\mathbb{C}}(\omega) = \overline{\omega}\}/\mathbb{R}^\times. \tag{1.1}$$

Note that every point in Ω_ψ is orthogonal to the lattice S.

Let $((X, \tau, \varphi),\ \alpha)$ be a marked real 2-elementary K3 surface of type (S, θ) satisfying $\alpha \circ \varphi_* \circ \alpha^{-1} = \psi$, i.e., "the associated integral involution" with $((X, \tau, \varphi,)\ \alpha)$ is ψ.

Let $H\ \ (\subset H_2(X, \mathbb{C}))$ be the Poincare dual of the complex 1-dimensional space $H^{2,0}(X)$. The 1-dimensional subspace $\alpha_{\mathbb{C}}(H)$ in $\mathbb{L}_{K3} \otimes \mathbb{C}$ is regarded as an element of Ω_ψ.

Definition 1.12. We say $\alpha_{\mathbb{C}}(H)$ the *period* of a marked real 2-elementary K3 surface $((X, \tau, \varphi),\ \alpha)$ of type (S, θ) satisfying $\alpha \circ \varphi_* \circ \alpha^{-1} = \psi$.

If we change their markings appropriately, the periods of all marked real 2-elementary K3 surfaces whose associated integral involutions are isometric to $(\mathbb{L}_{K3}, \psi)$ with respect to G can be found in the domain Ω_ψ. See [8].

Definition 1.13 (Equivalence)**.** We say a point $[v]$ $(\in \Omega_\psi)$ is *equivalent* to a point $[v']$ $(\in \Omega_\psi)$ if $[v'] = f_{\mathbb{C}}([v])$ for an automorphism f of $(\mathbb{L}_{K3}, \psi)$ of type (S, θ) with respect to the group G.

Lemma 1.14 ([8])**.** *If a point $[v]$ $(\in \Omega_\psi)$ is equivalent to $[v']$ $(\in \Omega_\psi)$ and $[v]$ is the period of some marked real 2-elementary K3 surface $((X, \tau, \varphi),\ \alpha)$ of type (S, θ) satisfying $\alpha \circ \varphi_* \circ \alpha^{-1} = \psi$, then $[v']$ is also the period of a marked real 2-elementary K3 surface $((X, \tau, \varphi),\ \alpha')$ of type (S, θ) satisfying $(\alpha') \circ \varphi_* \circ (\alpha')^{-1} = \psi$ where α' is some another marking of (X, τ, φ).*

Remark 1.15 ([6])**.** By the global Torelli theorem, if two periods are equivalent, then corresponding marked real 2-elementary K3 surfaces are analytically isomorphic (see Definition 1.11). The converse is also true.

The domain Ω_ψ has two connected components which are interchanged by $-\psi$. Since $-\psi$ is an automorphism of $(\mathbb{L}_{K3}, \psi)$ with respect to the group G, by Lemma 1.14 and Remark 1.15, it is enough to investigate the quotient space

$$\Omega_\psi / -\psi. \tag{1.2}$$

Now we set $\mathbb{L}_{\pm} := \{x \in \mathbb{L}_{K3} \mid \psi(x) = \pm x\}$.

We restrict ourselves the case $S \subset \mathbb{L}_-$, i.e., $\theta = -\mathrm{id}$, and we set

$$\mathbb{L}_{-,S} := \mathbb{L}_- \cap S^{\perp}.$$

For $[v] \in \Omega_\psi$ $(v \in \mathbb{L}_{K3} \otimes \mathbb{C})$, we consider the orthogonal decomposition $v = v_+ + v_-$ $(v_\pm \in \mathbb{L}_\pm \otimes \mathbb{R})$. Then we have $v_- \in \mathbb{L}_{-,S} \otimes \mathbb{R}$ and $v_+^2 = v_-^2 > 0$. Hence, both $\mathbb{L}_+$ and $\mathbb{L}_{-,S}$ are hyperbolic lattices.

Let $\mathcal{L}_+$, $\mathcal{L}_{-,S}$ be the hyperbolic spaces obtained from $\mathbb{L}_+ \otimes \mathbb{R}$, $\mathbb{L}_{-,S} \otimes \mathbb{R}$ respectively.

Then we have

$$\Omega_\psi / -\psi \;=\; \mathcal{L}_+ \times \mathcal{L}_{-,S} \quad \text{(a direct product)}. \tag{1.3}$$

1.4. $(\mathcal{D}\mathbb{R})$-nondegenerate marked real 2-elementary K3 surfaces.

Definition 1.16. An element $x(\neq 0) \in \mathrm{Pic}(X) \otimes \mathbb{R}$ is *nef* (for X) if $x \cdot C \geq 0$ for any effective curve C in $\mathrm{Pic}(X)$.

Definition 1.17 ([6])**.** We say a marked 2-elementary K3 surface $((X, \tau),\ \alpha)$ of type S is *$(\mathcal{D})$-degenerate* if there exists an element $x_0 \in \alpha_{\mathbb{R}}^{-1}(\mathcal{M})$ which is **not** nef.

This condition is equivalent to the existence of an exceptional curve (i.e., irreducible [1] and having negative self-intersection) with square -2 on the quotient surface $Y := X/\tau$. And this condition is also equivalent to have an element $\delta \in \mathrm{Pic}(X)$ with $\delta^2 = -2$ such that $\delta = (\delta_1 + \delta_2)/2$ where $\delta_1 \in S$, $\delta_2 \in S^{\perp}_{\mathrm{Pic}(X)}$, and $\delta_1^2 = \delta_2^2 = -4$.

We say a marked real 2-elementary K3 surface $((X, \tau, \varphi),\ \alpha)$ of type (S, θ) is *$(\mathcal{D}\mathbb{R})$-degenerate* if there exists a "real" element $x_0 \in \alpha_{\mathbb{R}}^{-1}(S_- \cap \mathcal{M})$ which is **not** nef, where we set

$$S_{\pm} := \{x \in S \,|\, \theta(x) = \pm x\}.$$

This condition is equivalent to have an element $\delta \in \mathrm{Pic}(X)$ with $\delta^2 = -2$ such that $\delta = (\delta_1 + \delta_2)/2$ where $\delta_1 \in S$, $\delta_2 \in S^{\perp}_{\mathrm{Pic}(X)}$ and $\delta_1^2 = \delta_2^2 = -4$, and δ_1 must be orthogonal to an element $x \in S_- \cap \mathrm{int}(\mathcal{M})$, where $\mathrm{int}(\mathcal{M})$ denote the interior part of $\mathcal{M}$.

Considering associated integral involutions, we get a natural map from the period domain of marked real 2-elementary K3 surfaces of type (S, θ) to the set of isometry classes with respect to G of integral involutions of $\mathbb{L}_{K3}$ of type (S, θ). We have

Theorem 1.18 ([6])**.** *The natural map above gives a bijective correspondence between the connected components of the period domain of $(\mathcal{D}\mathbb{R})$-nondegenerate marked real 2-elementary K3 surfaces of type (S, θ) and the isometry classes with respect to G of integral involutions of $\mathbb{L}_{K3}$ of type (S, θ) such that the fixed part $\mathbb{L}_{K3}^{\psi}$ of ψ is hyperbolic.*

2. Real 2-elementary K3 surfaces of type $((3, 1, 1), -\mathrm{id})$

Now we consider 2-elementary K3 surfaces of type S with the invariants

$$(r(S), a(S), \delta(S)) = (3, 1, 1).$$

We fix a sublattice S $(\cong (3, 1, 1))$ of the K3 lattice $\mathbb{L}_{K3}$.

We quote the following results from Alexeev and Nikulin [1]. See also [7] and [8].

Let (X, τ) be a 2-elementary K3 surface of type S $(\cong (3, 1, 1))$ and

$$A := X^{\tau}$$

[1] We say a class in $H_2(X, \mathbb{Z})$ is *irreducible* if it is represented by an irreducible divisor of X.

be the fixed point set (curve) of τ. Then we have

$$A = A_0 \cup A_1 \quad \text{(disjoint union)},$$

where A_0 is a nonsingular rational curve ($\cong \mathbb{P}^1$) with $A_0^2 = -2$ and A_1 is a nonsingular curve of genus 9. (X, τ) has a structure of an elliptic pencil $|E + F|$ with its exceptional section A_0 and the unique reducible fiber $E + F$ having the following properties:

(i): E is an irreducible nonsingular rational curve with $E^2 = -2$ and $E \cdot A_0 = 1$.

(ii): $E \cdot F = 2$, $F^2 = -2$, $F \cdot A_0 = 0$, and F is either
an irreducible nonsingular rational curve ("*type IIa case*"), or
the union of two irreducible nonsingular rational curves F' and F'' which are conjugate by τ, $F' \cdot F'' = 1$, and $(F')^2 = (F'')^2 = -2$ ("*type IIb case*"). See §2.4 of [1].

(iii): The classes $[A_0]$, $[E]$ and $[F]$ generate the lattice $H_{2+}(X, \mathbb{Z})$ ($\cong S$). Moreover, $A_1 \cdot E = 1$, $A_1 \cdot F = 2$. The Gram matrix of the lattice $H_{2+}(X, \mathbb{Z})$ with respect to the basis $[E]$, $[F]$ and $[A_0]$ is as follows:

	$[E]$	$[F]$	$[A_0]$
$[E]$	-2	2	1
$[F]$	2	-2	0
$[A_0]$	1	0	-2

Then we have an orthogonal decomposition

$$H_{2+}(X, \mathbb{Z}) = \mathbb{Z}([A_0],\ [E] + [F]) \oplus \mathbb{Z}([F]),$$

where the subgroups $\mathbb{Z}([A_0],\ [E] + [F])$ and $\mathbb{Z}([F])$ are isometric to the hyperbolic plane and $\langle -2 \rangle$ respectively.

For any real 2-elementary K3 surface (X, τ, φ) of type (S, θ) with $S \cong (3, 1, 1)$, we have

$$\theta = -\mathrm{id}$$

and

$$G = \{\mathrm{id}\},$$

where id stands for the identity map on S. See [7].

Remark that all real 2-elementary K3 surfaces of type $((3, 1, 1), -\mathrm{id})$ are $(\mathcal{D}\mathbb{R})$-non-degenerate.

We set $Y := X/\tau$ and let $\pi : X \to Y$ be the quotient map. We define the curves on Y as follows:

$$e := \pi(E) \quad \text{and} \quad f := \pi(F).$$

If F is the union of two nonsingular rational curves F' and F'' which are conjugate by τ and $F' \cdot F'' = 1$, then we have $f = \pi(F) = \pi(F' \cup F'') = \pi(F') = \pi(F'')$.

We use the same symbols A_0 and A_1 for their images in Y by π. Then, the Picard group $\mathrm{Pic}(Y)$ of Y is generated by the classes $[e]$, $[f]$ and $[A_0]$. The Gram matrix of $\mathrm{Pic}(Y)$ with respect to the basis $[e]$, $[f]$ and $[A_0]$ is as follows:

	$[e]$	$[f]$	$[A_0]$
$[e]$	-1	1	1
$[f]$	1	-1	0
$[A_0]$	1	0	-4

We have $A_1 \cdot e = 1$ and $A_1 \cdot f = 2$.

We now contract the exceptional curve $f = \pi(F)$ to a point and get a blow up

$$\mathrm{bl} : Y \to \mathbb{F}_4,$$

where $\mathbb{F}_4$ is the 4-th Hirzebruch surface.

We set $s := \mathrm{bl}(A_0)$, $A_1' := \mathrm{bl}(A_1)$, $c := \mathrm{bl}(e)$. Then s is the exceptional section of $\mathbb{F}_4$ with $s^2 = -4$ and c is a fiber of the fibration $\mathbb{F}_4 \to s$ with $c^2 = 0$.

We have

$$\mathrm{bl}(A) = \mathrm{bl}(A_0) + \mathrm{bl}(A_1) = s + A_1' \quad \in |-2K_{\mathbb{F}_4}|.$$

Namely, $\mathrm{bl}(A)$ is an anti-bicanonical curve. We have $-2K_{\mathbb{F}_4} \sim 12c + 4s$ and $A_1' \quad \in |12c + 3s|$.

Since $s \cdot A_1' = 0$, A_1' does not intersect the section s.

Remark 2.1 ([7], [8])**.** F is a nonsingular rational curve if and only if A_1 intersects f in two distinct points, and F is a union of two nonsingular rational curves if and only if A_1 touches f.

If A_1 intersects with f at two distinct points, then they are real points or non-real conjugate points. In the former case the real double point of A_1' is a real node, and in the latter case it is a real isolated point. Anyway the real double point of A_1' is nondegenerate. If A_1 touches to f in Y, then the real double point is a real cusp. (a degenerate double point)

Since all real 2-elementary K3 surfaces of type $((3,1,1), -\mathrm{id})$ are $(\mathcal{D}\mathbb{R})$-non-degenerate, the connected components of the period domain of marked real 2-elementary K3 surfaces of type $((3,1,1), -\mathrm{id})$ and the isometry classes with respect to $G = \{\mathrm{id}\}$ of integral involutions of $\mathbb{L}_{K3}$ of type $((3,1,1), -\mathrm{id})$ such that the fixed part $\mathbb{L}_{K3}^{\psi}$ of ψ is hyperbolic are in bijective correspondence by Theorem 1.18.

However, as is written in Remark 8 in Subsection 2.4 in [7], it is possible that an isometry class of integral involutions of $\mathbb{L}_{K3}$ of type $((3,1,1), -\mathrm{id})$ corresponds to **both type IIa case and type IIb case**. In other words, isometry classes cannot distinguish **degenerate and nondegenerate double points** of the curves A_1' on $\mathbb{F}_4$ (Recall Remark 2.1).

We would like to determine whether the real double point of the curve A_1' on $\mathbb{F}_4$ is degenerate or not.

To do this, we define a more strict marking α of a real 2-elementary K3 surface (X, τ, φ) of type $((3,1,1), -\mathrm{id})$. Compare Definition 2.2 below to Definition 1.4.

Let

$$\mathcal{E}, \quad \mathcal{F}, \quad \text{and} \quad \mathcal{A}$$

be generators of S with the Gram matrix

$$\begin{array}{c|ccc} & \mathcal{E} & \mathcal{F} & \mathcal{A} \\ \mathcal{E} & -2 & 2 & 1 \\ \mathcal{F} & 2 & -2 & 0 \\ \mathcal{A} & 1 & 0 & -2, \end{array}$$

$\mathcal{M} \cdot \mathcal{E} \geq 0$, $\mathcal{M} \cdot \mathcal{F} \geq 0$, and $\mathcal{M} \cdot \mathcal{A} \geq 0$.

We set

$$\mathbb{U} := \mathbb{Z}(\mathcal{A},\ \mathcal{E} + \mathcal{F}) \quad \text{and} \quad \langle -2 \rangle := \mathbb{Z}(\mathcal{F}).$$

Then $\mathbb{U}$ is isometric to the hyperbolic plane, and we have an orthogonal decomposition

$$S = \mathbb{U} \oplus \langle -2 \rangle.$$

Definition 2.2 (marked real 2-elementary K3 surfaces $((X, \tau, \varphi), \alpha)$ of type $((3,1,1), -\mathrm{id})$)**.** We define a *marked real 2-elementary K3 surfaces* $((X, \tau, \varphi), \alpha)$ *of type* $((3,1,1), -\mathrm{id})$ to be a pair of a real 2-elementary K3 surface (X, τ, φ) of type $((3,1,1), -\mathrm{id})$ (Definition 1.3) and a marking (isometry)

$$\alpha : H_2(X, \mathbb{Z}) \cong \mathbb{L}_{K3}$$

such that

- $\alpha(H_{2+}(X, \mathbb{Z})) = S$,

- $\alpha \circ \varphi_* = \theta \circ \alpha$ on $H_{2+}(X, \mathbb{Z})$,
- $\alpha_{\mathbb{R}}^{-1}(V^+(S))$ contains a hyperplane section of X,
- the set $\alpha^{-1}(\Delta(S)_+)$ contains only classes of effective curves of X,
- $\alpha([A_0]) = \mathcal{A}$, $\alpha([E]) = \mathcal{E}$, and $\alpha([F]) = \mathcal{F}$.

Note that every real 2-elementary K3 surface (X, τ, φ) of type $((3,1,1), -\mathrm{id})$ has such a marking α.

We give a criterion for the unique double point of a real anti-bicanonical curve $\mathrm{bl}(A) = s + A_1'$ (on $\mathbb{RF}_4$) with one real double point on A_1' to be nondegenerate.

Let us **fix** an integral involution

$$(\mathbb{L}_{K3}, \psi)$$

of type $((3,1,1), -\mathrm{id})$, and consider the period domain (Subsection 1.3)

$$\Omega_\psi / -\psi \ = \ \mathcal{L}_+ \times \mathcal{L}_{-,S} \ .$$

Lemma 2.3. *Let $[[\omega]]$ be a point [2] in $\Omega_\psi/-\psi$. Then, the real double point of the curve A_1' on $\mathbb{F}_4$ (Remark 2.1) which corresponds to $((X,\tau,\varphi),\alpha)$ is nondegenerate if and only if there are no $\mathbf{v}$ $(\neq \pm\mathcal{F})$ in $\mathbb{L}_{K3}$ satisfying that $\mathbf{v}\cdot\omega = 0$, $\mathbf{v}\cdot\mathbb{U} = 0$, and $\mathbf{v}^2 = -2$.*

Proof. The $(\Longleftarrow)$ direction has been proved in [8]. In this paper we prove the $(\Longrightarrow)$ direction. Assume that the double point of the real anti-bicanonical curve $\mathrm{bl}(A)$ on $\mathbb{F}_4$ is nondegenerate. Then F is irreducible ([7], [8]). Suppose that there **exists** a $\mathbf{v}$ $(\neq \pm\mathcal{F})$ in $\mathbb{L}_{K3}$ satisfying that $\mathbf{v}\cdot\omega = 0$, $\mathbf{v}\cdot\mathbb{U} = 0$, and $\mathbf{v}^2 = -2$. Then $-\mathbf{v}$ has the same properties. Hence, we may assume $\alpha^{-1}(\mathbf{v})$ is an effective class. $\tau^*(\alpha^{-1}(\mathbf{v}))$ is also effective. We have $\tau^*(\alpha^{-1}(\mathbf{v}))^2 = -2$, $\tau^*(\alpha^{-1}(\mathbf{v}))\cdot[A_0] = 0$, $\tau^*(\alpha^{-1}(\mathbf{v}))\cdot([E]+[F]) = 0$ and $\alpha(\tau^*(\alpha^{-1}(\mathbf{v})))\cdot\omega = 0$ since τ^* is non-symplectic. Hence, $\alpha(\tau^*(\alpha^{-1}(\mathbf{v})))$ has the same properties as $\mathbf{v}$. Since $\alpha(\tau^*(\alpha^{-1}(\mathbf{v}))) + \mathbf{v} \in S$, it is contained in $\mathbb{Z}(\mathcal{F})$. There exists an integer n such that

$$\alpha(\tau^*(\alpha^{-1}(\mathbf{v}))) + \mathbf{v} = n\mathcal{F}.$$

Let us consider a hyperplane section class $h \in \mathrm{Pic}\, X$, which is ample. Since $\tau^*(\alpha^{-1}(\mathbf{v}))$ and $\alpha^{-1}(\mathbf{v})$ are effective, we have $\tau^*(\alpha^{-1}(\mathbf{v}))\cdot h > 0$ and $\alpha^{-1}(\mathbf{v})\cdot h > 0$. We have $\tau^*(\alpha^{-1}(\mathbf{v})) + \alpha^{-1}(\mathbf{v}) \neq 0$. Hence, we have $n \neq 0$. Moreover, since $\tau^*(\alpha^{-1}(\mathbf{v})) + \alpha^{-1}(\mathbf{v})$ and $[F](= \alpha^{-1}(\mathcal{F}))$ are effective, we have $n > 0$. Actually, if $n < 0$, then $0 = (\tau^*(\alpha^{-1}(\mathbf{v})) + \alpha^{-1}(\mathbf{v}) + (-n)\alpha^{-1}(\mathcal{F}))\cdot h > 0$. This is a contradiction.

If $\tau^*(\alpha^{-1}(\mathbf{v})) = \alpha^{-1}(\mathbf{v})$, then $(\tau^*(\alpha^{-1}(\mathbf{v})) + \alpha^{-1}(\mathbf{v}))^2 = (2\alpha^{-1}(\mathbf{v}))^2 = -8$. On the other hand, $(\tau^*(\alpha^{-1}(\mathbf{v})) + \alpha^{-1}(\mathbf{v}))^2 = (n[F])^2 = -2n^2$. Hence, $n = 2$. Thus we have $\mathbf{v} = \mathcal{F}$. This is a contradiction. Thus we have $\tau^*(\alpha^{-1}(\mathbf{v})) \neq \alpha^{-1}(\mathbf{v})$.

Let $\alpha^{-1}(\mathbf{v}) = \sum_i m_i \mathbf{v}_i$, where m_i are positive integers and $\mathbf{v}_i$ are irreducible effective classes. Obviously we have $\alpha(\mathbf{v}_i)\cdot\omega = 0$ for any i.

If $\mathbf{v}_i = [E]$ or $[F]$, then $\mathbf{v}_i\cdot([E]+[F]) = 0$. And if $\mathbf{v}_i \neq [E]$ and $\neq [F]$, then we have $\mathbf{v}_i\cdot([E]+[F]) \geq 0$ for any i. Here E is irreducible, and moreover, F is irreducible by the assumption. Since $\alpha^{-1}(\mathbf{v})\cdot([E]+[F]) = 0$, we have $\mathbf{v}_i\cdot([E]+[F]) = 0$ for any i.

Since $[A_0]\cdot([E]+[F]) = 1$, we see $\mathbf{v}_i \neq [A_0]$ for any i. Thus we also have $\mathbf{v}_i\cdot[A_0] \geq 0$. Since $\alpha^{-1}(\mathbf{v})\cdot[A_0] = 0$, we have $\mathbf{v}_i\cdot[A_0] = 0$ for any i.

Thus, we have $\alpha(\mathbf{v}_i)\cdot\mathbb{U} = 0$ for any i. Since $\mathbb{U}$ is of signature $(1,1)$, we have $(\mathbf{v}_i)^2 < 0$ for any i by the Hodge index theorem. Such $\mathbf{v}_i$'s are rational $(\cong \mathbb{P}^1)$ and we have $\mathbf{v}_i^2 = -2$.

[2] By the surjectivity of the period map of marked K3 surfaces ([2], p.339), $[\omega]$ is the period of a marked real 2-elementary K3 surface $((X,\tau,\varphi),\alpha)$ of type $((3,1,1),-\mathrm{id})$ satisfying $\alpha\circ\varphi_*\circ\alpha^{-1} = \psi$.

Suppose $\alpha(\mathbf{v}_i) = \mathcal{F}$ for some i, say $i = 1$. Namely, $\mathbf{v}_1 = [F]$. Let $\mathbf{v}' := \alpha^{-1}(\mathbf{v}) - m_1[F]$. Since $\tau^*(\alpha^{-1}(\mathbf{v})) \neq \alpha^{-1}(\mathbf{v})$ (see above), we have $\mathbf{v}' \neq 0$. Thus, there exists a $\mathbf{v}_i (\neq \pm[F])$ in $H_2(X, \mathbb{Z})$ such that $\mathbf{v}_i \cong \mathbb{P}^1$ (irreducible), $\mathbf{v}_i^2 = -2$, $\alpha(\mathbf{v}_i) \cdot \omega = 0$, and $\alpha(\mathbf{v}_i) \cdot \mathbb{U} = 0$.

Eventually we can choose $\mathbf{v}$ such that $\alpha^{-1}(\mathbf{v})$ is an irreducible class.

Since $\tau^*(\alpha^{-1}(\mathbf{v})) \neq \alpha^{-1}(\mathbf{v})$, $\tau^*(\alpha^{-1}(\mathbf{v}))$ and $\alpha^{-1}(\mathbf{v})$ are represented by different irreducible curves respectively. Hence, we have $\tau^*(\alpha^{-1}(\mathbf{v})) \cdot \alpha^{-1}(\mathbf{v}) \geq 0$.

Since $(\alpha(\tau^*(\alpha^{-1}(\mathbf{v}))) + \mathbf{v})^2 = (-2) + (-2) + 2(\tau^*(\alpha^{-1}(\mathbf{v})) \cdot \alpha^{-1}(\mathbf{v})) = -2n^2$, we have $2 - n^2 \geq 0$. Thus, we have $n = 1$. Namely, we have

$$\mathcal{F} = \mathbf{v} + \alpha(\tau^*(\alpha^{-1}(\mathbf{v}))),$$

namely,

$$[F] = \alpha^{-1}(\mathbf{v}) + \tau^*(\alpha^{-1}(\mathbf{v})).$$

Let F' be an irreducible curve representing $\alpha^{-1}(\mathbf{v})$, and F'' be $\tau^*(F')$. Thus $F' + F''$ represents the class $[F]$.

Hence, there exists a marked real 2-elementary K3 surface corresponding to the period $[\omega]$ which has $E + F' + F''$ as a unique reducible fiber of its elliptic fibration.

Conversely, every marked real 2-elementary K3 surface corresponding to the period $[\omega]$ has $E + F' + F''$ as a unique reducible fiber of its elliptic fibration.

Hence, we have $F = F' + F''$. This contradicts the assumption that F is irreducible. This completes the proof of Lemma 2.3. □

Acknowledgments. The author would like to thank Professor Hisanori Ohashi for his helpful suggestions and comments.

References

[1] V.A. Alexeev, V.V. Nikulin, *Del Pezzo and K3 Surfaces,* MSJ Memoirs, 2006.

[2] W.P.Barth, K. Hulek, C.A.M.Peters, and A.Van de Ven, *Compact Complex Surfaces,* Springer, 2004.

[3] I. Itenberg, *Curves of degree* 6 *with one non-degenerate double point and groups of monodromy of non-singular curves,* Real Algebraic Geometry, Proceedings, Rennes 1991, Lecture Notes in Math., **1524**, Springer, (1992), 267–288.

[4] V.V. Nikulin, *Integral symmetric bilinear forms and some of their geometric applications,* Math. USSR Izv., **14** (1980), 103–167.

[5] V.V. Nikulin, *On the quotient groups of the automorphism groups of hyperbolic forms by the subgroups generated by* 2*-reflections, Algebraic-geometric applications,* J. Soviet Math., **22** (1983), 1401–1476.

[6] V.V. Nikulin, Sachiko Saito, *Real K3 surfaces with non-symplectic involution and applications,* Proc. London Math. Soc., **90** (2005), 591–654.

[7] V.V. Nikulin, Sachiko Saito, *Real K3 surfaces with non-symplectic involution and applications. II,* Proc. London Math. Soc., **95** (2007), 20–48.

[8] Sachiko Saito, *On real anti-bicanonical curves with one double point on the 4-th real Hirzebruch surface,* Journal of Singularities **11** (2015), 1–32.

Department of Mathematics Education, Asahikawa Campus,, Hokkaido University of Education, Asahikawa, JAPAN

E-mail address: saito.sachiko@a.hokkyodai.ac.jp